



09876980-060802




Approximated Peano Curve. The space-filling process has not been completed.

Figure 1A (Prior Art)




Boustrophedon Path

Figure 1B (Prior Art)



Archimedes Spiral defined by equally distributed points

Figure 1C (Prior Art)



Spiral-like line-based scanning

Figure 1D (Prior Art)




Figure 2A




Figure 2B




Figure 3A




Figure 3B

09876980.060801




Figure 3C






Figure 3D

0876980.060801




The situation of Lemma 1

Figure 4A




Case (A)

Figure 4B




Case (B)

Figure 4C




Smooth transition between two circles of different radii.

Figure 4D



Smooth transition between two circles of same radius.


Figure 4E



Transition between two unit circles of radius 1. The distance between the circles is $\sqrt{3}$


Figure 4F

09876980-060801



Beginning (left) and completion (right) of a scanning scheme where the curvature is below a certain value


Figure 5A



Construction of $s2$ and the subsequent part of the curve

Figure 5B

09876980.060801



Conformal Spiral.

Figure 6

09876980.060801






Figure 07

09876980.060801



The first 1000 Halton points (left) and randomly chosen points (right)

Figure 8A



Original solution (left) and splined version (right).

Figure 8B

09376980-060801





Figure 9



Definition of I_1 , I_2 , and I_3

Figure 10




Figure 11A




Figure 11B




Figure 11C




Figure 12A






Figure 12B

09876980.060801



Beginning of a Low Discrepancy Curve based on a specific Halton Sequence in 2d


Figure 13A



Splined Low Discrepancy Curve coarse search


Figure 13B

108090-08692860



Comparison of Conformal Spiral and Low Discrepancy Searching

Figure 13C



Comparison of Travel Distance for Low Discrepancy Search and Conformal Spiral Search

Figure 13D

09876980-060801






Figure 14




Tiling of the plane and relation to the surface of the unit cube



Low-discrepancy curve in a ring


Figure 15A

Figure 15B




Low Discrepancy Preserving Mapping Function

Figure 15C



Low-discrepancy curve filling the surface of a torus

Figure 15D



Low-discrepancy curve on a sphere
(left) and projections (right)


Figure 16


```

graph TD
    1702[Select a Parameterization of a Surface S Where the Parameter Space is the Unit Square  
1702] --> 1704[Select a LD Curve in the Parameter Space, i.e.,  
Select a Suitable (alpha1, alpha2)  
1704]
    1704 --> 1706[Determine a Re-Parameterization of S Such That the Ratio of Line and Area Elements of S, Based on a Riemannian Metric, is Constant  
1706]
    1706 --> 1708[Map the LD Curve in the Unit Square onto the Surface S Using the Re-Parameterization  
1708]
    1708 --> 1710[Output the Mapped LD Curve on S  
1710]

```


Figure 17



Surfaces can be scanned efficiently when the term low discrepancy sequence/ curve can be generalized, e.g. based on metrics on the surface.

Figure 18

098765980.060801




Splined Low Discrepancy Curve coarse search with refined final approach

Figure 19

08876980-060801


Intensity Field Distribution in Search Area



Beam intensity distribution in search area


Figure 20

09876980.060801





Location of the Peak

Figure 21A



Initial Final Approach Move

Figure 21B



Error distribution of the estimated peak X coordinate error (left) and Y coordinate error (right)

Figure 21C




Figure 22




Figure 23